

EFFECT OF WEED CONTROL METHODS ON THE PERFORMANCE OF SELECTED VARIETIES OF MAIZE (*ZEAMAYS* L.) AT JALINGO AND GOMBE, NORTH-EASTERN NIGERIA

¹Garba, M.D, ²Bakari Aliyu and ²Chukudi G. Michael

¹Department of Agricultural Science Education, Federal College of Education (Technical), Gombe and ²Taraba State University, Jalingo

Tel: 08036473176, 08061402640, email: manugarba082@gmail.com

ABSTRACT

A field experiment was conducted at Jalingo (Taraba State) and Gombe (Gombe State), Nigeria, during 2023 and 2024 raining season. In Jalingo the experiment was conducted at the Teaching and Research Farm, Taraba State University and Dadin Kowa area in Gombe. This study was conducted to evaluate the effect of weed control methods on the performance of selected varieties of Maize (*Zea mays* L.). To assess weed flora composition and population dynamics in maize fields at 3, 6, and 9 weeks after sowing (WAS). Weed species were identified and grouped into broadleaf, grasses, and sedges. The results demonstrate marked spatial variation in weed community structure and successional trends, emphasizing the need for location-specific weed management strategies in maize production systems of northeastern Nigeria. The experimental design was a split-plot design (SPD) replicated three times. The total experimental area was, 26m x 73m and plot size was 3m x 3.5m with a distance of 2m between replicates and 1m between plots, respectively. Maize varieties :V1 (=Sammaz 29), V2(= Sammaz 34), V3 (= Sammaz 53), V4 (=Sammaz 54), V5 (= Tzer White) and V6 (=Tzer Yellow (QPM)) were allocated to the main plot and while weed control methods application rates I used 150 ml per knapsack sprayer (PRE (=Alachlor, POE(= Glyphosate) at kg/ha,PPO(= Paraquats at kg/ha, SPT (=Sweet potato), RHW (=Recommended hand weeding) and UWC (= Unweeded control) at kg/ha, were randomly assigned to subplot replicated three times. Data collected includes plant height, leaf area, fresh grass weight, 100 seed weight and grain yield (Kg/ha). Data obtained were subjected to analysis of variance (ANOVA) using SAS software and their means were separated using DMRT. Results obtained revealed significant variations in maize plant parameters, due to maize variety, weed control methods. V1 (Sammaz 29 under mean combined Jalingo and Gombe over the two (2) years identified significant performance than other variety in term of growth and yield which recorded (2676.8Kg/ha) whereas PRE treatment indicated the highest with (2803.3Kg/ha) and the least goes to UWC (2229.5Kg/ha). Therefore Variety V1 observed the highest grain yields whereas V6 indicated the tallest plant height. However V1, V6, PRE, POE and RHW are recommended for cultivation and weed control at both Jalingo and Gombe.

Keywords: Maize variety, Weed flora, population dynamics, savanna ecology, Jalingo, Gombe

INTRODUCTION

Maize (*Zea mays* L.) belongs to the grass family poaceae and is believed to have originated from Mexico or Central America and spread to West Africa by early European traders in the 16th century (Saleem, 2022). It is the world's third most important crop after rice (*Oryza sativa* L.) and wheat (*Triticum aestivum* L.). A total of 951.6 million metric tonnes of maize are produced worldwide, with 1.5 tonnes/ha in Africa, 1.7 tonnes/ha in India, slightly more than tonnes/ha in Latin America, 12.5 tonnes/ha in the USA and in Nigeria, with an annual production of 11.3 tonnes/ha (FAOSTAT, 2023).

Maize is the most important cereal crop in Sub-Saharan Africa (SSA) as documented by Dauda and Ahmed, (2024). It is high yielding, easy to process, readily digested and cheaper than other cereals. Every part of maize plant has economic value; the grain, leaves, stalk, tassel and cob can all be used to produce a large variety of food and non- food products . In industrialized countries, maize is largely used as livestock feed and as raw material for industrial products (Bates, 2022), while in developing countries it is mainly used for human consumption and poultry feed (Luis and Reigner, 2024).

Green maize (fresh on the cob) is eaten parched, baked, roasted or boiled, playing an important role in filling the hunger gap after the dry season (Saidu and Isa, 2022). Maize production in West and Central Africa has witnessed a phenomenal increase in the last two decades, the level of production was low before as at 2010, but the level has seriously increased (Goswami *et al.*, 2023). Maize grain contains about 70% carbohydrate, 10% protein, 4% oil, 2.3% crude fibre, 10.4% albuminoides and 1.4% ash depending on the variety (Bukar and David, 2022). The grain has significant quantities of vitamin A, nicotinic acid, riboflavin, Vitamin B, vitamin C, ascorbic acid, vitamin E. It is low in calcium and fairly high in phosphorous (Baquest *et al.*, 2023).

MATERIALS AND METHODS

The experiments were carried out in two locations Jalingo and Gombe in North- Eastern Nigeria. Jalingo field trial was conducted at Taraba State University Teaching and Research Farm in the Department of Agronomy Jalingo is located in the Northern Guinea Savanna ecological zone of Nigeria and lies between Lat. 8°38'00"N; Long 10°46'00"E). Jalingo has annual rainfall ranges from 800 - 1,063.2mm from the months of April to October (Adebayo and Orunoye, 2022). Gombe field trial was conducted at Dadin Kowa (Lat. 10°15'N; Long 11°15'E. Gombe is located in the Northern Guinea Savannah ecological zone of Nigeria. Gombe has annual rainfall ranges from 850 - 1000mm from the months of May to October (Kwari *et al.*, 2022). The experimental materials consisted of six varieties of maize:-V1 (SAMMAZ 29); V2 (SAMMAZ 34); V3 (SAMMAZ 53); V4 (SAMMAZ 54); V5 (TZER-WHITE) and V6 (TZER-YELLOW) Quality protein maize (QPM). These varieties have varying degree of yield potential maturity period, drought tolerant, and striga tolerance resistance and were obtained from the Institute for Agricultural Research (IAR) Samaru, Zaria, Kaduna state, (first four) and the remaining two varieties from Federal University Kashere, Gombe state Nigeria. The seeds from both sources were certified seeds; sweet potatoes vines were obtained at Kasuwan Bera while the six weed control methods were PRE, POE, PPO, SPT, RHW and UWC.

The experimental design used was a split of plot design (SPD) in randomized complete block design replicated three times. The total experimental area was 26m x 73m. Each plot size was 3m x 3.5m. The varieties were randomly assigned to each replication as main plot. The weed control options were randomly assigned within each main plot. Five

plants per plot were randomly tagged for periodic observations during crop growth stages, at 3, 6 and 9 WAS. Maize growth parameters assessed were; plant height, number of leaves, leaf area and fresh grass weight. Yield and yield components measured were; Maize above ground biomass, 100 seed weight, and yield grain (kg/ha). Data collected were subjected to analysis of variance (ANOVA) using SAS software and their means were separated using DMRT.

RESULTS AND DISCUSSIONS

Effect of Maize Varieties and Weed control methods on Plant height of maize were computed and analyzed for combined location in Jalingo and Gombe within the 2023 and 2024 Raining Season. The results of the Varieties (V1-V6) and Weed control methods (PRE - UWC), are presented in Table 1. Jalingo at Weed control methods RHW gave the tallest (123.68cm) and the shortest was observed in UWC. Gombe at 3, 6 and 9WAP, V6 identified better than the others in term of plant height, Therefore, at Weed control methods (WCM), PRE recorded better (126.22cm) and the least goes to RHW.

Table 2. Jalingo at 3WAP 2024, the shortest plant height is pointed out in V1(18.01), therefore at Weed Control Methods (WCM), the shortest was observed in PRE. At 6WAP, V6 gave the tallest and the shortest is recorded in V1(32.31cm). Under Weed Control Methods (WCM), PRE observed better.

At 9WAP, V4 shown the tallest (128.04cm) and the shortest indicated in V2 (117.50cm). However, at Weed Control Methods, PPO recorded better height while the shortest indicated in UWC (113.68cm).

Gombe at 3WAP, the varietal effect shown that V2 observed better, whereas at Weed Control Methods, PRE shown better (19.17cm). At 6WAP, V5 and V4 identified better than the rest. At Weed Control Methods, SPT and POE observed better than the rest.

At 9WAP, V5 is better (160.28cm). At Weed Control Methods, PRE (167.97CM and shortest goes to POE (148.37cm), respectively.

Average across location at 3WAP, V2 shown better. Therefore, at Weed control methods, which is highly significant.

Average across location, at 6WAP, the shortest indicated in V1 (41.07cm). However, at Weed Control Methods, SPT shown better. Average across location, at 9WAP, the varietal effect V5 observed better than the rest. Therefore, at Weed Control Methods, RHW is the least (18.38cm).

Table 1: Effect of Variety and Weed Control Methods on Plant height (PH) (cm) of Maize at Jalingo and Gombe in 2023 Raining Season.

Means with same letters are not significantly differences from each other according to DMRT

Variety	Jalingo			Gombe			Mean		
	3	6	9	3	6	9	3	6	9
V1	16.68b	34.14c	104.03b	17.97b	40.74b	117.73ab	17.32c	37.44d	110.88b
V2	18.50ab	42.01b	106.67b	19.30ab	44.10b	115.35b	18.90a-c	43.05bc	111.01b
V3	19.86a	44.26b	112.23b	19.94ab	45.63b	115.14b	9.90ab	44.94b	113.68b
V4	19.20a	39.91bc	126.21ab	18.81ab	38.98b	110.72b	19.02ab	39.45cd	118.47b
V5	18.33ab	41.45b	104.94b	18.44ab	39.41b	114.56b	18.39bc	40.43b-d	109.75b
V6	20.32a	52.23a	144.48a	20.72a	54.09a	134.23a	20.52a	53.16a	139.36a
WEED CONTROL METHODS									
PRE	19.23ab	50.0 0ab	113.40a	19.44a	43.53a	126.22a	19.34ab	42.26ab	119.81a
POE	19.70a	47.01a	116.18a	19.68a	46.90a	115.27a	19.69ab	46.99a	115.73a
PPO	16.93b	38.93b	114.31a	18.24a	40.94a	110.78a	17.59c	39.94b	112.54a
SPT	18.25ab	41.88ab	117.96a	18.38a	43.54a	121.01a	18.31bc	42.71ab	119.49a
RHW	18.75ab	41.39ab	123.68a	19.37a	44.63a	123.09a	19.06a-c	43.01ab	123.38a
UWC	20.07a	43.72ab	113.03a	20.07a	43.42a	111.36a	20.07a	43.57ab	112.20a
VxWCT	ns	Ns	Ns	Ns	*	Ns	***	ns	Ns

at $p \leq 0.05$ V1=Sammaz 29, V2=Sammaz 34, V3= Sammaz 53, V4=Sammaz 54, V5=Tzer White and V6=Tzer Yellow (QPM), PRE = Alachlor, POE = Glyphosate, PPO = Paraquats, SPT = Sweet potato (Smother Crop), RHW = Recommended hand weeded, UWC = Unweeded control, WCM = Weed control methods* WCM = Variety verse weed control methods and WAP =Weeks after planting.

Table 2: Effect of Variety and Weed Control Methods on Plant height (PH) (cm) at Jalingo and Gombe 2024 Raining Season

Variety	Jalingo Mean			Gombe					
	3	6	9	3	6	9	3	6	9
V1	18.01a	32.31d	118.53b	18.12a	49.84a	160.18a	18.07a	41.07b	139.35a
V2	18.35a	40.34bc	117.50b	19.08a	51.11a	158.34a	18.72a	45.72ab	137.34a
V3	18.06a	42.64b	121.48ab	18.79a	50.85a	159.91a	18.42a	46.74ab	140.69a
V4	18.34a	35.03cd	128.04a	19.03a	53.21a	146.39a	18.69a	44.12ab	137.22a
V5	18.36a	39.89bc	126.67ab	18.84a	54.52a	160.28a	18.60a	47.20ab	142.10a
V6	18.30a	49.65a	125.12ab	18.84a	47.93a	154.08a	18.57a	48.79a	139.19a
WEED CONTROL METHODS									
PRE	17.86a	40.43a	121.77ab	19.17a	50.38a	167.97a	18.52a	45.41a	144.87a
POE	18.95a	39.99a	125.83a	18.60a	51.57a	148.37b	18.77a	45.78a	136.44a
PPO	18.07a	39.82a	127.01a	18.92a	50.23a	158.02ab	18.49a	45.03a	142.52a
SPT	18.08a	40.28a	122.17a	18.44a	52.91a	157.67ab	18.26a	46.60a	139.92a
RHW	18.01A	39.68a	126.88a	18.83a	51.13a	153.73ab	18.42a	45.41a	18.38ab
UWC	18.45a	39.65a	113.68b	18.75a	51.22a	152.17ab	18.60a	45.44a	19.23a
VxWCT	ns	Ns	Ns	ns	*	ns	***	ns	ns

Means with same letters are not significantly differences from each other according to DMRT at $p \leq 0.05$ V1=Sammaz 29, V2=Sammaz 34, V3= Sammaz 53, V4=Sammaz 54, V5=Tzer White and V6=Tzer Yellow (QPM), PRE = Alachlor, POE = Glyphosate, PPO = Paraquats, SPT = Sweet potato (Smother Crop), RHW = Recommended hand weeded, UWC = Unweeded control, WCM = Weed control methods* WCM = Variety verse weed control methods and WAP =Weeks after planting.

Effect of Variety and Weed Control Methods on Leave Area (LA) cm^2 at Jalingo and Gombe 2023 Raining Season

Effect of variety and weed control methods on leaf area of maize were computed and analyzed at Jalingo and Gombe location. The results as presented in Table 3 showed that for Varieties V1-V6 and Weed control methods PRE - UWC. Jalingo at 3, 6 and 9WAP, the least leaf area indicated in V5 (1465.6cm^2). However at Weed Control Methods, PRE Plots gave the higher and the least is recorded in PPO (285.11cm^2). At Gombe in 3, 6 and 9WAP, V6 is better, while the least goes to V2 (1356.2cm^2). Therefore at Weed Control Methods, PPO gave the highest (1763.6cm^2).

The results as presented in Table 4, showed that for Varieties V1-V6 and Weed control methods PRE - UWC.

At Jalingo 2024, at 3WAP, V5 (105.93cm^2) is better and the least indicated in V2 (86.09cm^2). At Weed Control Methods, POE performed better. At 6WAP, V2 (197.98cm^2) shown better them the rest of the varieties. At Weed Control Methods, the least recorded in

POE Plots (179.10cm²). At 9WAP, V6 (665.5cm²) recorded better, under Weed Control Methods, indicated RHW (671.9cm²).

Gombe 2024, at 3WAP, V1 (107.84cm²) shown better and Weed Control Methods, UWC (104.88cm²) shown better performance. At 6WAP, the least identified in V2 (161.58cm²). However at Weed Control Methods SPT recorded better (190.25cm²). At 9WAP, V5 (505.80cm²) indentified better Variety whereas Weed Control Methods, PPO (505.62cm²) is better than the rest of varieties.

Average across location at 3WAP, the Varieties recorded the same performance ,but V1 is better whereas the least identified in V2 (90.55cm²). Therefore at Weed Control Methods, POE shown better. At 6WAP, Average across location, V3 (187.94cm²) identified better under Weed Control Methods showing similar performance, but RHW observed different.

Average across location at 9WAP, the Varietal effect observed the same pattern, while V6 (577.35cm²) also at Weed Control Methods, RHW (592.71cm²) observed better in performance.

Table 3: Effect of Variety and Weed Control Methods on Leave Area (LA) cm² at Jalingo and Gombe 2023 Raining Season

	Jalingo			Gombe			Mean		
Vari	3	6	9	3	6	9	3	6	9
V1	342.95a	1672.0	6206.	434.83a	1684.6a	6708.	388.89a	1678.3a	6457.9a
V2	282.09a	1483.6	6392.	331.00a	1356.2b	6320.	306.55a	1419.9c	6356.8b
V3	340.58a	1522.3	6811.	443.60a	1543.3a	6592.	392.09a	1532.8b	6702.3a
V4	362.90a	1648.6	6811.	356.02aab	1539.8a	7092.	359.46a	1594.2b	6951.8a
V5	332.62a	1465.6	6251.	388.82ab	1722.3a	6371.	360.72a	1594.0c	6311.6b
V6	407.88a	1968.8	6861.	378.30ab	1804.0a	7344.	393.09a	1886.4a	7102.6a
WEED CONTROL METHOD									
PRE	477.77a	1644.7	6798.	402.12a	1529.5a	6772.	439.94a	1587.1a	6785.1a
POE	329.47a	1784.0	6434.	430.07a	1430.3a	6693.	379.77a	1607.2a	6563.8a
PPO	285.11b	1474.2	6307.	333.10a	1763.6a	7249.	309.10b	1618.9a	6778.2a
SPT	358.05a	1588.6	6362.	361.48a	1597.7a	6553.	359.76a	1593.2a	6457.7a
RH	286.71b	1545.4	6404.	395.75a	1625.1a	6323.	341.23a	1585.2a	6363.7a
UW	331.92a	1724.0	7029.	410.06a	1704.0a	6838.	370.99a	1714.0a	6934.3a
VxW	Ns	*	Ns	Ns	Ns	Ns	ns	Ns	ns

Means with same letters are not significantly differences from each other according to DMRT at $p \leq 0.05$ V1=Sammaz 29, V2=Sammaz 34, V3= Sammaz 53, V4=Sammaz 54, V5=Tzer White and V6=Tzer Yellow (QPM), PRE = Alachlor, POE = Glyphosate, PPO = Paraquats, SPT = Sweet potato (Smother Crop), RHW = Recommended hand weeded, UWC = Unweeded control, WCM = Weed control methods* WCM = Variety verse weed control methods and WAP =Weeks after planting.

Table 4: Effect of Variety and Weed Control Methods on Leave Area (cm²) of maize at Jalingo and Gombe 2024 Raining Season

Variety	Jalingo			Gombe			Mean		
	3	6	9	3	6	9	3	6	9
V1	104.31a	176.65a	430.1a	107.84a	189.71a	504.99a	106.07a	183.18a	467.54a
V2	86.09b	197.98a	438.5a	95.00a	161.58b	499.95a	90.55b	179.78a	468.34a
V3	101.02ab	189.72a	446.8a	105.44a	186.15a	508.25a	103.23ab	187.94a	477.54a
V4	99.91ab	195.52a	411.7a	100.77a	171.52ab	488.98a	100.34ab	183.52a	450.34a
V5	105.93a	183.14a	444.4a	92.73a	190.15a	505.80a	99.33ab	186.64a	474.24a
V6	96.21ab	183.67a	665.5a	93.43a	184.04a	483.98a	94.86ab	183.86a	577.35a
WEED CONTROL METHOD									
PRE	104.98a	189.12ab	431.3a	99.02a	178.79a	497.18	101.10a	183.96a	464.23
POE	105.47a	179.10b	426.5a	98.86a	179.34a	501.33a	102.26a	179.22a	461.73a
PPO	92.28a	184.63ab	423.8a	103.62a	182.38a	505.62a	97.95a	183.51a	464.72a
SPT	95.96a	184.14ab	444.6a	100.10a	190.25a	499.34a	98.03a	187.20a	471.95a
RHW	99.01a	202.86a	671.9a	89.04a	182.11a	508.84	94.03a	192.48a	592.71a
UWC	95.77a	186.82ab	438.9a	104.88a	170.29a	480.87	100.32a	178.56a	459.90a
VxWCT	ns	Ns	ns	ns	ns	ns	ns	Ns	ns

at $p \leq 0.05$ V1=Sammaz 29, V2=Sammaz 34, V3= Sammaz 53, V4=Sammaz 54, V5=Tzer White and V6=Tzer Yellow (QPM), PRE = Alachlor, POE = Glyphosate, PPO = Paraquats, SPT = Sweet potato (Smother Crop), RHW = Recommended hand weeded, UWC = Unweeded control, WCM = Weed control methods* WCM = Variety verse weed control methods and WAP =Weeks after planting.

Effect of between Variety and Weed Control Methods on Fresh Grass Weigh (FGW) (g/m²) at Jalingo and Gombe location 2023 and 2024 Raining Season

Effect of Variety and Weed control methods on Fresh Grass Weigh g/m² (FGW) g/m² were computed and analyzed, at Jalingo and Gombe location 2023 and 2024. The results as presented in Table 5, showed that Varieties V1-V6 and Weed control methods PRE - UWC. At Jalingo in 2023, V2 (24.70 g/m²) recorded better performance than the rest. However at weed control methods, UWC observed the heaviest with (38.02 g/m²). At Gombe 2023, V1 (23.71 g/m²) whereas at Weed Control Methods, the least identified in RHW (13.93g/m²). Average across location V2 identified better than the other varieties. Therefore at Weed

Control Methods, UWC observed the heaviest whereas the least indicated in RHW (12.69g/m²).

Jalingo 2024, only V4 is far better than the rest of the varieties which indicated (110.08g/m²). However at Weed Control Methods, UWC recorded the heaviest Fresh Grass Weight of Grass (135.22g/m²) and the least goes to POE (33.39g/m²). At Gombe 2024, only V6 shown the heaviest then the rest. Therefore at Weed Control Methods, UWC (263.46 g/m²) gave the heaviest while the least indicated in SPT (35.04g/m²).

Table 5: Effect of between Variety and Weed Control Methods on Fresh Grass Weigh (FGW) (g/m²) at Jalingo and Gombe location 2023 and 2024 Raining Season

	2023			2024			Mean (Combine)		
Variety	Jalingo	Gombe	Mean	Jalingo	Gombe	Mean	Jalingo	Gombe	Mean
V1	16.85 a	23.71a	20.28 a	38.69a	55.86a	47.27a	27.77a	39.78a	33.78a
V2	24.70 a	22.47a	23.59 a	35.91a	132.76a	82.95a	30.30a	76.04a	52.85a
V3	22.07 a	20.07a	21.07 a	41.48a	48.17a	44.83a	31.78a	34.12a	32.95a
V4	16.93 a	21.63a	19.28 a	110.08a	61.97a	86.03a	63.50a	41.80a	52.65a
V5	18.05 a	21.57a	19.81 a	44.21a	38.56	41.47a	31.13a	29.83a	30.49a
V6	16.32 a	21.85a	19.09 a	38.87a	133.44a	84.80a	27.59a	76.05a	51.48a
WEED CONTROL METHOD									
PRE	20.01 b	18.54b	19.28 b	34.73b	43.19b	38.96b	27.37b	30.87b	29.12b
POE	13.85 b	19.23b	16.54 b	33.39b	36.23b	34.73b	23.62b	27.23b	25.37b
PPO	11.36 b	17.74b	14.55 b	34.28b	40.43b	37.36b	22.82b	29.09b	25.95b
SPT	20.24 b	17.44b	18.84 b	37.91b	35.04b	36.47b	29.07b	26.24b	27.65b
RHW	11.45 b	13.93b	12.69 b	33.71b	41.84b	37.66b	22.58b	27.49b	25.00b
UWC	38.02 a	44.43a	41.22 a	135.22a	263.46a	199.34a	86.62a	153.94a	120.28a
VxWCT	Ns	**	***	Ns	ns	ns	ns	Ns	ns

Means with same letters are not significantly differences from each other according to DMRT at $p \leq 0.05$ V1=Sammaz 29, V2=Sammaz 34, V3= Sammaz 53, V4=Sammaz 54, V5=Tzer White and V6=Tzer Yellow (QPM), PRE = Alachlor, POE = Glyphosate, PPO = Paraquats, SPT = Sweet potato (Smother Crop), RHW = Recommended hand weeded, UWC = Unweeded control, WCM = Weed control methods* WCM = Variety verse weed control methods and WAP =Weeks after planting.

Effect of Variety and Weed Control Methods on Maize Above Ground Biomass (MAGB) (g/m²) Yield of Maize five (5) sample Plants at Jalingo and Gombe location 2023 and 2024 Raining Season

The results are presented in Table 6, showed that for Varieties V1-V6 and Weed control methods PRE - UWC. At Jalingo 2023, V5 observed the heaviest than the rest with (127.93 g/m²) and least identified in V2 (118.86 g/m²). However at Weed Control Methods, the heaviest observed in RHW plot with (131.01g/m²), whereas the least was recorded in UWC (116,23 g/m²).

Gombe 2023, the Varietal effect identified similar trend, while V6 indicated the heaviest (132.54 g/m²). Therefore at Weed Control Methods, PPO recorded the heaviest with (131.63g/m²). Average across location, the Varietal effect observed in the same vine, but V3 gave the heaviest (125.68 g/m²), therefore, at Weed Control Methods, RHW gave the heaviest (127.61g/m²) and the least pointed out in UWC (108.08 g/m²).

Jalingo 2024, Varietal effect and Weed control methods showed drastically reduction in their growth yield performance compared with Gombe location, because Jalingo in the year 2024 experience longer period of drought. V4 recorded better than the rest with (99.44 g/m²) and the least indicated in V5 (81.03 g/m²), whereas under Weed control methods, SPT recorded better performance with (99.82g/m²). Gombe 2024, V1 recorded the heaviest among the rest with (502.03 g/m²). However, at Weed Control Methods; which indicated similar direction, whereas UWC observed the highest performance (318.12 g/m²).

Table 6: Effect of Variety and Weed Control Methods on Above Ground Biomass (AGB) (g/m²) Yield of Maize five (5) sample Plants at Jalingo and Gombe location 2023 and 2024 Raining Season

	2023			2024			(Combine)		
Variety	Jalingo	Gombe	Mean	Jalingo	Gombe	Mean	Jalingo	Gombe	Mean
V1	122.68a	119.35a	121.01a	86.82a	502.03a	294.43a	104.75a	310.69a	207.72a
V2	118.86a	112.44a	115.65a	84.32a	481.66a	277.32a	101.59a	291.78a	195.35a
V3	125.41a	125.96a	125.68a	85.20a	460.40a	272.80a	105.30a	293.18a	199.24a
V4	126.11a	119.37a	122.74a	99.44a	426.78a	263.11a	112.77a	273.08a	192.93a
V5	127.93a	107.14a	117.53a	81.03a	468.35a	269.16a	104.48a	282.58a	192.28a
V6	118.97a	132.54a	125.56a	85.90a	494.40a	284.31a	102.43a	313.47a	204.94a
WEED CONTROL METHOD									
PRE	125.01ab	128.92a	126.96a	88.74a	512.21a	300.48a	106.87a	320.57a	213.72a
POE	124.26ab	115.04ab	119.65ab	91.46a	491.07a	279.51a	107.86a	292.00a	197.30a
PPO	120.69ab	131.63a	126.16a	90.55a	496.44a	293.49a	105.62a	314.04a	209.83a
SPT	122.75ab	116.15ab	119.55ab	99.82a	497.35a	298.58a	111.28a	312.20a	210.33a
RHW	131.01a	124.21ab	127.61a	92.77a	447.62a	265.13a	111.89a	281.30a	195.40a
UWC	116.23b	99.92b	108.08b	59.38b	318.12a	223.75a	87.81b	244.02a	165.91a
VxWCT	ns	Ns	ns	Ns	ns	ns	ns	ns	ns

Means with same letters are not significantly differences from each other according to DMRT at $p \leq 0.05$ V1=Sammaz 29, V2=Sammaz 34, V3= Sammaz 53, V4=Sammaz 54, V5=Tzer White and V6=Tzer Yellow (QPM), PRE = Alachlor, POE = Glyphosate, PPO = Paraquats, SPT = Sweet potato (Smother Crop), RHW = Recommended hand weeded, UWC = Unweeded control, WCM = Weed control methods* WCM = Variety verse weed control methods and WAP =Weeks after planting.

Effect of Variety and Weed Control Methods on 100 Seed Weight at Jalingo and Gombe location 2023 and 2024 Raining Season

Effect of Variety and Weed control methods on 100 Seed Weight were computed and analyzed at Jalingo and Gombe location 2023 and 2024. The result are presented in Table 7 showed that for Varieties V1-V6 and Weed control methods PRE - UWC. At Jalingo 2023, the Varietal effect identified that V4 recorded the highest weight among others with (20.60g), however at Weed Control Methods, PRE observed the highest with (20.68g), while the least goes to POE with (19.22g).

Gombe 2023, the Varieties observed similar performance, but V5 indicated the highest (19.29g) whereas the least recorded in V3 with (17.08g). However at Weed Control Methods, UWC recorded the highest with (19.77g). Average across location the Varietal effect observed that V1 recorded better than the rest with (19.67g), therefore at Weed control methods, PRE pointed out the heaviest (19.65g). Jalingo 2024, V6 identified the heaviest among others (20.28g) and at Weed Control Methods, RHW recorded the highest (20.58g) the least goes to UWC (18.72g/m²). Gombe V6 is shown better performance than the rest (24.66g), however at Weed Control Methods, UWC gave (24.54g). Jalingo and Gombe over two (2) years, the Varietal effect were highly significant throughout the experiment, V1 recorded the highest number of 100 seed weight with (20.99g), therefore at Weed Control Methods, RHW identified more effective than the others with (20.80g), respectively.

Table 7: Effect of Variety and Weed Control Methods on 100 Seed Weight at Jalingo and Gombe location 2023 and 2024 Raining Season

	2023			2024			Mean (Combine)		
Variety	Jalingo	Gombe	Mean	Jalingo	Gombe	Mean	Jalingo	Gombe	Mean
V1	20.54a	18.80 a	19.67a	19.99a	24.65a	22.32a	20.27a	21.72a	20.99a
V2	18.68a	18.39 a	18.53a	19.84a	23.90ab	21.81a	19.26a	21.07a	20.15a
V3	19.96a	17.08 a	18.52a	19.30a	23.53ab	21.42a	19.63a	20.31a	19.97a
V4	20.60a	17.83 a	19.21a	18.94a	24.42a	21.68a	19.77a	21.12a	20.45a
V5	19.29a	19.29 a	19.29a	19.25a	21.95b	20.56a	19.27a	20.59a	19.92a
V6	20.52a	18.04 a	19.28a	20.28a	24.66a	22.41a	20.40a	21.26a	20.82a
WEED CONTROL METHOD									
PRE	20.68ab	18.63 ab	19.65a	19.79ab	22.84a	21.32a	20.23ab	20.74a	20.49a
POE	19.22b	18.14 ab	18.68a	19.75ab	24.04a	21.77a	19.49a	20.92a	20.18a
PPO	21.66a	16.42 b	19.04a	18.92ab	23.62a	21.27a	20.29ab	20.02a	20.15a
SPT	18.39b	18.22 ab	18.30a	19.85ab	24.01a	21.93a	19.12ab	21.11a	20.12a
RHW	20.41ab	18.26 ab	19.33a	20.58a	24.17a	22.32a	20.49a	21.13a	20.80a
UWC	19.24b	19.77 a	19.50a	18.72b	24.54a	21.63a	18.98b	22.16a	20.57a
VxWCT	*	ns	ns	*	***	ns	***	ns	ns

Means with same letters are not significantly differences from each other according to DMRT at $p \leq 0.05$ V1=Sammaz 29, V2=Sammaz 34, V3= Sammaz 53, V4=Sammaz 54, V5=Tzer White and V6=Tzer Yellow (QPM), PRE = Alachlor, POE = Glyphosate, PPO = Paraquats, SPT = Sweet potato (Smother Crop), RHW = Recommended hand weeded, UWC = Unweeded control, WCM = Weed control methods* WCM = Variety verse weed control methods and WAP =Weeks after planting.

Effect of Variety and Weed Control Methods on Grain Yield (Kg/ha) at Jalingo and Gombe in 2023 and 2024 Raining Season

Effect of Variety and Weed control methods on grain yield kg/ha were computed and analyzed at Jalingo and Gombe location 2023 and 2024. The results are presented in Table 8 showed that for Varieties V1-V6 and Weed control methods PRE - UWC.

At Jalingo in 2023 the Varietal effect were significant but V3 recorded the highest Maize Grain Yield (MGY) with (2320.8 Kg/ha) and the least indicated in V5 plot (1624.4 Kg/ha). However at Weed control methods, PPO observed the highest (2686.9kg/ha) and the least goes to POE (1651.7kg/ha).

At Gombe the Varietal effect shown that V1 is far better than the other varieties with (2336.4 Kg/ha) but the least identified at V3 (1540. 6Kg/ha, therefore under weed control methods, PRE recorded the highest (2498.2kg/ha) the least goes to SPT (1631.4kg/ha). Average across location in 2023 are the same in Varietal effect, whereas V1 gave the highest performance with (2315.4kg/ha) and the least pointed out in V5 (1901.5kg/ha. However at Weed control methods, PRE pointed out the highest (2530.6kg/ha).

Table 8: Effect of Variety and Weed Control Methods on Grain Yield (Kg/ha) at Jalingo and Gombe in 2023 and 2024 Raining Season

	2023			2024			Mean (Combine)		
Variety	Jalingo	Gombe	Mean	Jalingo	Gombe	Mean	Jalingo	Gombe	Mean
V1	2294.5a	2336.4a	2315.4a	1774.6a	4301.7a	3038.1a	2034.5a	3319.0a	2676.8a
V2	1701.5a	2159.5ab	1930.5a	1492.2a	4068.7a	2780.5a	1596.9a	3114.1a	2355.5a
V3	2320.8a	1540.6b	1930.7a	1512.5a	3825.7a	2669.1a	1916.6a	2683.2a	2299.9a
V4	2163.4a	2116.7ab	2140.1a	1640.5a	4277.0a	2958.7a	1901.9a	3196.9a	2549.4a
V5	1624.4a	2178.6ab	1901.5a	1844.2a	3511.8a	2678.0a	1734.3a	2845.2a	2289.8a
V6	2270.7a	1949.4ab	2110.0a	1653.5a	4118.3a	2885.9a	1962.1a	3033.8a	2498.0a
WEED CONTROL METHODS									
PRE	2563.1ab	2498.2a	2530.6a	1757.6a	4394.5a	3076.0a	2160.3a	3446.3a	2803.3a
POE	1651.7c	1858.3ab	1755.0c	1821.0a	3883.4a	2852.2a	1736.4ab	2870.8a	2303.6ab
PPO	2686.9a	1918.5ab	2302.7ab	1512.6ab	3719.9a	2616.2a	2099.7a	2819.2a	2459.5ab
SPT	1825.3c	1631.4b	1728.4c	2057.2a	4373.3a	3215.3a	1941.3a	3002.4a	2471.8ab
RHW	1936.8bc	2326.6ab	2131.7abc	1628.9ab	3714.3a	2671.6a	1782.8ab	3020.4a	2401.6ab
UWC	1711.5c	2048.4ab	1879.9bc	1140.1b	4017.8a	2579.0a	1426.8b	3033.1a	2229.5b
VxWCT	ns	ns	*	ns	ns	ns	*	ns	ns

Means with same letters are not significantly differences from each other according to DMRT at $p \leq 0.05$ V1=Sammaz 29, V2=Sammaz 34, V3= Sammaz 53, V4=Sammaz 54, V5=Tzer White and V6=Tzer Yellow (QPM), PRE = Alachlor, POE = Glyphosate, PPO = Paraquats, SPT = Sweet potato (Smother Crop), RHW = Recommended hand weeded, UWC = Unweeded control, WCM = Weed control methods* WCM = Variety verse weed control methods and WAP =Weeks after planting.

CONCLUSION

The variation observed between the two locations underscore the strong influence of environmental conditions on weed community composition. Overall, effective weed management in maize must consider species-specific dominance patterns and seasonal succession to minimize yield losses.

RECOMMENDATIONS

Variety V1 (SAMMAZ 29) for maize grain yield (MGY) and Variety V6 (Tzer yellow) (QPM) for the tallest plant height is hereby recommended in the study areas for Varieties stated.

Farmers in both Jalingo and Gombe should adopt PRE, POE and RHW for the weed control method.

REFERENCES

- Adebayo, V. and Orunoye T. (2023): Optimum dose of nitrogen for higher yielding and dwarf maize. *African Journal of crop science* 16(11):12-17.
- Baquest, M, Evan, J, Melvin, C and Victor, A. (2023). Cropping system sub-programmed. In farming system Research programme. Institute for Agricultural Research Ahmadu Bello University Zaria. Nigeria pp 112-118.
- Bates, Y.(2022). Threatening Food self-sufficiency:GATT; impact on the grains industry in the GATT: Philippine Issues and perspectives: Philinine cereal and tuber Institute, Quezon City.
- Bukar, H. and David, O (2022): Striga infestation in Kenya status distribution and management options sustained agriculture. *Research programme* 2: 99-105.
- Dauda, S and Ahmed. N.(2024). Review on maize based intercropping. *Nigerian Journal of Agronomy* 9: 135-147..
- FAOSTAT, (2023).): Ecological interpretation of weed flora dynamics under different tillage systems. *Agriculture Ecosystem and Environment*. 66:177-183.
- Goswami, N.N, Kamath, M.B and Samatoso, D, Luis, V.R. (2023): Phosphorous requirement and management of maize, sorghum and wheat, In: Phosphorus requirement for sustainable agriculture in Asia trans Oceania. Pp 340-348.
- Kwari, C.O, Kimball, B.A, Ahuja, L.R. Pandey, P.O and Rayar, K.V. (2022):Simulation of maize evapotranspiration an inter-comparison among 29 maize models. *Agric for meteorology*: 271: 264-271 [Google scholar Cross Ref]
- Luis, S. and Reigner, T. (2024): A new strain of downy mildew pathogen of maize: A Lynne Rienner, Boulder.industry. In byerlee D., Eicher, C.K. (eds). *Africa's Emerging maize revolution*. Lynne Rienner, Boulder.
- Saidu, G and Isa, M. (2022). Agronomic and environmental implications of organic farming pattern. *African Journal of plant physiology* 8:543-550.
- Saleem. N.P. (2022): Effects of maize and legume intercropping systems and nitrogen on crop yields and residual nitrogen. *Field crops Research*, 4: 133-145.