

Climate Variability and Soil Moisture Dynamics in the South-South Region of Nigeria

Idisi E. B.^{1*}, Deekor T.N.², & Lawal, O.³

¹Department of Environmental Management, Faculty of Environmental Sciences, Delta State University of Science and Technology, Ozoro, Delta State, Nigeria.

²Department of Geography And Environmental Management, Faculty of Social Sciences, University of Port Harcourt, Rivers State, Nigeria.

³Department of Geography And Environmental Management, Faculty of Social Sciences, University of Port Harcourt, Rivers State, Nigeria.

*Corresponding Author: benjamin.idisi86@gmail.com

Abstract

This study examines the interplay between climate variability and soil moisture dynamics in the South-South region of Nigeria, focusing on their impacts on vegetation and agricultural productivity. Using long-term temperature, precipitation, and soil moisture datasets from 2000 to 2020, the research identifies significant trends and anomalies through statistical methods such as the Mann-Kendall trend test and Sen's slope estimator. Cross-wavelet analysis was employed to explore the relationships among climate variables and their combined influence on vegetation dynamics. The findings reveal a clear link between soil moisture variability and climatic conditions, emphasizing the critical role of soil moisture in moderating vegetation response to changes in precipitation and temperature. The study highlights the seasonal nature of soil moisture dynamics, with higher moisture retention during wet seasons supporting robust vegetation growth and lower levels during dry seasons exacerbating vegetation stress. It also identifies feedback mechanisms between soil moisture and atmospheric conditions, where variations in soil moisture influence local precipitation patterns and vegetation resilience. Furthermore, the impact of extreme weather events, such as floods and droughts, underscores the need for adaptive strategies to mitigate their effects on agriculture and water resources. Recommendations include promoting sustainable agricultural practices, enhancing soil moisture retention through agroforestry and improved irrigation, strengthening early warning systems for extreme weather events, and adopting remote sensing technologies for monitoring and resource management. By addressing the challenges posed by climate variability, these strategies aim to enhance resilience and support sustainable development in the region. This research provides a comprehensive framework for understanding the interconnections between climate, soil moisture, and vegetation, offering valuable insights for policymakers and stakeholders to develop targeted interventions in the face of a changing climate.

Keywords: Climate variability, Soil moisture dynamics, Vegetation response, Sustainable agriculture, Temporal Analysis

1.0 INTRODUCTION

Climate variability significantly influences soil moisture dynamics, particularly in regions like South-South Nigeria, where the interplay of climatic factors and soil characteristics can lead to pronounced effects on agricultural productivity and ecosystem health. Soil moisture is a critical component of the hydrological cycle, acting as a mediator between precipitation and vegetation health, and its variability is influenced by a multitude of factors including temperature, land cover, and atmospheric conditions (Roux *et al.*, 2013; Ghajarnia *et al.*, 2020). In the context of South-South Nigeria, understanding these dynamics is essential for effective land management and agricultural practices, especially given the region's vulnerability to climate change. The spatio-temporal variability of soil moisture is a well-documented phenomenon, shaped by local climatic conditions, soil heterogeneity, and vegetation cover (Ghajarnia *et al.*, 2020; Mittelbach and Seneviratne, 2012; Ekness and Randhir, 2015). In South-South Nigeria, the region's diverse topography and climatic conditions contribute to significant variations in soil moisture levels. For instance, areas with high vegetation cover tend to retain moisture better than those with sparse vegetation, highlighting the importance of land cover in moisture dynamics (Schwingshackl *et al.*, 2018; Myeni *et al.*, 2021). Moreover, the seasonal patterns of rainfall in the region, characterized by distinct wet and dry seasons, further complicate the soil moisture dynamics, necessitating a nuanced understanding of these interactions for effective agricultural planning (Vezhapparambu *et al.*, 2019; Lorenz *et al.*, 2016; Uguru *et al.*, 2022).

Research has shown that soil moisture plays a pivotal role in regulating surface temperature and precipitation patterns, particularly in tropical regions (Zhang *et al.*, 2011; Ardilouze *et al.*, 2018; Carón *et al.*, 2014). In the South-South region, the coupling between soil moisture and atmospheric conditions can lead to feedback loops that exacerbate climate variability. For example, increased soil moisture can enhance local evaporation rates, which in turn can influence cloud formation and precipitation patterns, creating a complex interplay that can lead to extreme weather events (Wing *et al.*, 2016; Berg *et al.*, 2014; Salley *et al.*, 2016). This feedback mechanism is particularly relevant in the context of climate change, as shifts in temperature and precipitation patterns are likely to alter soil moisture dynamics, with potential implications for agricultural productivity and water resource management (Song *et al.*, 2019; Li *et al.*, 2019; Wang *et al.*, 2018). The role of soil moisture in the hydrological cycle cannot be overstated; it is a key variable that influences not only local ecosystems but also broader climatic patterns (Krakauer *et al.*, 2010; Zhu *et al.*, 2019; Liu, 2023).

In South-South Nigeria, where agriculture is a primary livelihood, understanding the relationship between soil moisture and crop yield is critical. Studies have indicated that fluctuations in soil moisture can significantly impact crop growth and yield, particularly in regions that experience irregular rainfall patterns (Das and Maity, 2014; Saeed *et al.*, 2013; Han *et al.*, 2020). Furthermore, the persistence of soil moisture anomalies can affect seasonal weather forecasts, complicating agricultural planning and resource allocation (Li *et al.*, 2019; Senanayake *et al.*, 2022). In addition to its direct effects on vegetation and agriculture, soil moisture also interacts with groundwater levels and surface runoff, influencing the overall hydrological balance in the region (Ghajarnia *et al.*, 2020; Agrawal & Chakraborty, 2020; Fernandez-Long *et al.*, 2020). The interplay between soil moisture and these hydrological components is particularly crucial in the context of flood and drought management, as changes in soil moisture can lead to significant variations in runoff and groundwater recharge (Ekness & Randhir, 2015; Urban *et al.*, 2015; Morrison & Wagner, 2020). In South-South Nigeria, where flooding is a recurrent issue, understanding these dynamics is essential for developing effective flood management strategies.

The integration of remote sensing technologies has enhanced our ability to monitor soil moisture dynamics over large spatial scales, providing valuable data for understanding climate variability (Song *et al.*, 2021; Blank *et al.*, 2023). In South-South Nigeria, satellite-derived soil moisture data can complement ground-based observations, offering insights into the spatial distribution and temporal changes in soil moisture across different land cover types and climatic conditions (Huang *et al.*, 2022; Guo *et al.*, 2016). This technological advancement is particularly important for regions with limited ground-based monitoring infrastructure, as it enables researchers and policymakers to make informed decisions regarding land and water management. Moreover, the implications of climate change on soil moisture dynamics in South-South Nigeria are profound. Projections indicate that rising temperatures and altered precipitation patterns will likely exacerbate soil moisture variability, leading to increased incidences of drought and flooding. Understanding these trends is critical for developing adaptive strategies that can mitigate the impacts of climate change on agriculture and natural ecosystems in the region. For instance, implementing sustainable land management practices that enhance soil moisture retention can help buffer against the adverse effects of climate variability (Li and Rodell, 2013; Dirmeyer and Norton, 2018).

2.0 MATERIALS AND METHODS

2.1 Data Sources

The study of soil moisture dynamics and climate analysis in South-South Nigeria necessitates the integration of various data sources to ensure comprehensive and accurate results. Primary data sources include meteorological stations, satellite imagery, and hydrological models. Meteorological data, particularly rainfall and temperature records, are crucial for understanding climatic conditions and their impact on soil moisture. The Climate Hazards Group Infrared Precipitation with Station data (CHIRPS) and other satellite-based rainfall products have been evaluated for their performance in Nigeria, demonstrating varying degrees of accuracy and reliability in capturing precipitation patterns across different regions (Ogbuet *et al.*, 2020; Agbonaye and Izinyon, 2021; Usman *et al.*, 2018). These satellite datasets provide a spatially distributed view of rainfall, which is essential for hydrological modeling in data-sparse regions like South-South Nigeria. In addition to rainfall data, temperature records are vital for modeling evapotranspiration and understanding the thermal dynamics affecting soil moisture. The integration of temperature data from multiple sources, including ground stations and satellite observations, enhances the robustness of climate analysis (Ogbu *et al.*, 2020). Furthermore, soil moisture data can be derived from remote sensing techniques, which allow for the monitoring of soil moisture at various depths and spatial resolutions, providing insights into moisture availability for agricultural practices (Odusanya *et al.*, 2019; Okeola *et al.*, 2023). Geographical Information Systems (GIS) play a pivotal role in managing and analyzing spatial data. By integrating topographic, hydrological, and land use data, GIS facilitates the assessment of how these factors influence soil moisture and hydrological processes in the region (Adeogun *et al.*, 2023; Adeogun *et al.*, 2014). The use of GIS in conjunction with hydrological models, such as the Soil and Water Assessment Tool (SWAT), allows for the simulation of water balance components and the evaluation of management strategies for sustainable water resource use (Odusanya *et al.*, 2019; Adeogun *et al.*, 2023; Adeogun *et al.*, 2014). The study employed a combination of primary and secondary data. Satellite-based remote sensing products, such as MODIS (Moderate Resolution Imaging Spectroradiometer) and Landsat imagery, were utilized to extract vegetation health indicators, including the Normalized Difference Vegetation Index (NDVI). Climatic data, including precipitation, air temperature, and soil moisture levels, were sourced from global databases such as the ERA5 reanalysis dataset and NASA's POWER Climate

Data. Ground truthing and in-situ measurements were conducted at selected locations to validate remotely sensed data and ensure accuracy.

2.2 Soil Moisture Modeling

Soil moisture dynamics were modeled using data derived from satellite-based soil moisture products like SMAP (Soil Moisture Active Passive) and AMSR-E (Advanced Microwave Scanning Radiometer). The Temperature-Vegetation Dryness Index (TVDI) was applied to assess moisture variability across different ecological zones. Correlation and regression analyses were performed to determine the relationships between soil moisture and vegetation health indicators, incorporating local factors such as soil type and land use patterns. Soil moisture modeling is a critical component of hydrological studies, particularly in regions susceptible to climate variability and extreme weather events. Studies have demonstrated the effectiveness of satellite-based evapotranspiration data in calibrating SWAT models, enhancing the model's predictive capabilities in data-sparse regions (Odusanya *et al.*, 2019; Sholagberu, 2016; Adeogun *et al.*, 2023). Additionally, the use of remote sensing data allows for the continuous monitoring of soil moisture, providing real-time insights into moisture availability and its implications for agricultural productivity and water management (Odusanya *et al.*, 2019; Okeola *et al.*, 2023). In South-South Nigeria, the modeling of soil moisture must consider the unique climatic and geographical characteristics of the region (Figure 1). The interplay between rainfall patterns, temperature fluctuations, and soil properties significantly influences moisture retention and availability. For instance, the seasonal variability of rainfall and the impact of land use changes on soil moisture dynamics are critical factors that need to be integrated into the modeling framework (Merem *et al.*, 2019; Agbonaye and Izinyon, 2021; Adeogun *et al.*, 2023).




Figure 1: Soil Types in South-south Region of Nigeria

2.3 Climate Analysis

Climate variability in the South-South region of Nigeria was assessed using long-term temperature and precipitation data from 2000 to 2020. Temporal trends were analyzed using the Mann-Kendall trend test and Sen's slope estimator, while seasonal and annual anomalies were mapped to identify shifts in climatic patterns. Cross-wavelet analysis was applied to investigate the interactions between soil moisture, precipitation, and temperature, emphasizing their combined influence on vegetation dynamics. This approach offers a robust framework for examining the relationships between climate variables and soil moisture, providing critical insights into ecological responses to changing climatic conditions. The study of climate trends in the South-South region involves analyzing long-term temperature and precipitation data to understand their effects on soil moisture and agricultural practices. Historical climate datasets are examined to detect significant trends, which are essential for assessing the region's susceptibility to climate change (Merem *et al.*, 2019; Agbonaye and Izinyon, 2021). For instance, analyses of rainfall variability reveal notable fluctuations that can influence agricultural output and water resource management (Usman *et al.*, 2018). Statistical tools and climate models, such as General Circulation Models (GCMs), have been employed to project future climatic conditions, offering insights into how potential changes in precipitation patterns may affect water availability and hydrological processes (Dembélé *et al.*, 2022). Additionally, examining extreme weather events, including floods and droughts, is critical for formulating adaptive strategies to mitigate their impacts on agriculture and water resources. Identifying trends in extreme precipitation and their relationship with soil moisture can help inform policymakers and stakeholders about the need for enhanced water management and infrastructure development (Merem *et al.*, 2019). These efforts are vital for ensuring sustainable agricultural practices and resource management in the face of climate variability.

2.4 Soil and Vegetation

The South-South region of Nigeria, consisting of six states—Akwa Ibom, Bayelsa, Cross River, Delta, Edo, and Rivers—is distinguished by its unique combination of soil types and vegetation, shaped by the area's climate and geographic features. The region's humid tropical climate supports a rich array of plant and animal life, making it one of the country's ecologically significant zones (Uguru *et al.*, 2024). Soils in this area are primarily alluvial and hydromorphic, with varying fertility levels influenced by proximity to water sources such as the Niger River and the Atlantic Ocean. These soils play a vital role in the region's agricultural economy. The dominant vegetation in the South-South includes tropical rainforests, which gradually transition into mangrove forests along the coast. The rainforest is highly biodiverse, providing habitat for various tree species, shrubs, and wildlife. Mangrove forests, particularly prominent in Bayelsa and Rivers States, are critical for coastal protection and serve as breeding grounds for numerous marine species. The region's wetlands further contribute to its ecological wealth, supporting biodiversity and sustaining local fisheries (Enuoh and Ogogo, 2018). High organic matter from dense vegetation contributes to the generally fertile soils of the South-South region. However, agricultural activities, especially the cultivation of oil palm, cassava, and rice, have led to soil depletion in some areas. Environmental damage from oil exploration and production, including soil contamination from spills and gas flaring, has further degraded the ecosystem, threatening both agricultural productivity and environmental sustainability (Adesina *et al.*, 2020; Enuoh and Ogogo, 2018). Agriculture in the region includes both subsistence and commercial farming, with mixed cropping systems combining legumes, tubers, cereals, and cash crops. These practices support food security and economic resilience. However, the use of traditional farming

methods and challenges like flooding and erosion—exacerbated by climate change—raise concerns about the long-term sustainability of agriculture in the region (Tinubu *et al.*, 2019).

Efforts to address vegetation degradation have increasingly focused on sustainable practices, such as agroforestry and reforestation, aimed at restoring degraded lands and preserving biodiversity. Combining indigenous ecological knowledge with modern agricultural techniques is being explored as a way to ensure sustainable land management in the region. The South-South's ecological diversity also makes it a priority for conservation efforts. Programs targeting biodiversity protection, particularly in protected areas like Cross River National Park, aim to balance environmental preservation with the livelihoods of local communities (Enuoh and Ogogo, 2018). In summary, the South-South region's ecological and agricultural landscape is shaped by its climatic and human factors. While the area boasts abundant biodiversity and fertile soils that drive agriculture, it faces pressing environmental challenges from oil activities and unsustainable land use. Ensuring the ecological sustainability of the region will require adopting sustainable agricultural methods and conservation strategies that harmonize environmental protection with community development

3.0 RESULTS AND DISCUSSION

3.1 Air Temperature

In 2000, the minimum temperature recorded was 21.87°C, while the maximum reached 27.59°C. This baseline indicates relatively moderate temperatures for the region, with a difference of 5.72°C between the minimum and maximum (Figure 2). This year sets a reference point for tracking temperature changes in the following years.




Figure 2: Mean Air Temperature Across States in the South - south region in 2000

By 2003, there was a slight rise in both the minimum and maximum temperatures, with the minimum increasing to 22.17°C and the maximum to 27.77°C. The rise of 0.30°C in the

minimum temperature suggests warmer nights or cooler periods, while the increase in the maximum of 0.18°C points to slightly hotter days. These changes suggest a gradual warming trend, which aligns with regional and global climate patterns. In 2006, the minimum temperature increased to 22.28°C , and the maximum reached 27.99°C . Both the minimum and maximum temperatures continued their upward trend, with the maximum rising more significantly. This increase reflects a continued warming of the region, with more pronounced temperature extremes during the hottest parts of the day. The narrowing range between the minimum and maximum temperatures suggests more stable but slightly hotter conditions overall. In 2009, the minimum temperature slightly decreased to 22.26°C , while the maximum dropped to 27.84°C (Figure 3). The small decrease in the maximum temperature might reflect short-term climatic variations, such as increased cloud cover or rainfall, leading to less extreme daytime heat. However, the minimum temperature remained stable, suggesting that nighttime or cooler period temperatures were still high, maintaining the overall warming trend.




Figure 3: Mean Air Temperature Across States in the South - south region in 2009

A noticeable change occurred in 2012, with the minimum temperature dropping to 21.89°C and the maximum to 27.47°C . The decline in both values indicates a cooler year compared to the preceding period. This shift may have been influenced by local climatic conditions such as increased rainfall or a temporary cooling event. Despite the overall warming trend, this year stands out as an anomaly with lower temperature extremes. In 2015, temperatures began to rise again, with the minimum temperature increasing to 22.19°C and the maximum to 27.86°C . This recovery from the cooler 2012 values suggests the resumption of the warming trend seen earlier in the decade. The return to higher minimum and maximum temperatures indicates a consistent warming trajectory, likely driven by factors such as deforestation and urbanization in the region. The year 2017 marked a significant increase in both minimum and maximum temperatures. The minimum temperature rose to 22.48°C , while the maximum reached 28.21°C (Figure 4). This year recorded some of the highest temperature extremes, with the maximum exceeding 28°C for the first time. The increased maximum suggests hotter days, while the higher minimum points to warmer nights or cooler periods, indicating that overall temperatures were higher across all times of day. In 2020, the South-South region

recorded its highest minimum and maximum temperatures over the two decades, with the minimum reaching 22.57°C and the maximum climbing to 28.38°C. The maximum temperature in 2020 is the highest in the entire dataset, signifying more intense heat during the day. The rise in the minimum temperature to 22.57°C also suggests that nights or cooler periods were warmer than in previous years. This increase in both minimum and maximum temperatures underscores the region's vulnerability to ongoing global warming, with potential impacts on agriculture, health, and local ecosystems.




Figure 4: Mean Air Temperature Across States in the South - south region in 2017

3.2 Precipitation

2000: Moderate Minimum and Maximum Precipitation

In 2000, the South-South geopolitical zone experienced a minimum daily rainfall of 1.45 mm and a maximum of 5.39 mm. The gap between these values suggests moderate variation in rainfall patterns during this year (Figure 5). The minimum rainfall indicates a relatively low level of precipitation on certain days, while the maximum value of 5.39 mm signifies significant rainfall on peak days. This range reflects a generally balanced distribution of rainfall, where neither extremely low nor very high precipitation events dominated the region during this period. By 2003, both the minimum and maximum daily rainfall values saw a slight decline, with the minimum recorded at 1.04 mm and the maximum at 5.22 mm. The reduction in both parameters suggests a decrease in the overall intensity of rainfall compared to 2000. Although the difference between the minimum and maximum values remained similar, the lowered figures point to a year characterized by less extreme rainfall events. This could indicate milder weather conditions and a lesser risk of both droughts and flooding. In 2006, there was an increase in the maximum daily rainfall to 5.82 mm, while the minimum value rose slightly to 1.32 mm (Figure 6). This indicates a moderate increase in rainfall intensity, with higher peak rainfall days. The maximum value of 5.82 mm suggests more significant rain events than in previous years, potentially leading to localized flooding or heavy rain days. The slight increase in the minimum value, however, still suggests that some days experienced low rainfall, maintaining variability in precipitation.




Figure 5: Mean Precipitation Across States in the South - south region in 2000




Figure 6: Mean Precipitation Across States in the South - south region in 2006

The year 2009 saw a significant jump in maximum rainfall, reaching 6.88 mm, while the minimum stayed relatively stable at 1.35 mm. This substantial increase in maximum rainfall points to the occurrence of heavier rain events, which may have had implications for flooding risks and water resource management in the region. The minimum value remaining stable suggests that while peak rainfall days became much more intense, lower precipitation days did not change significantly, preserving variability. In 2012, there was an interesting shift, with the minimum daily rainfall rising to 1.55 mm, the highest recorded in the 20-year period, while the maximum rainfall dropped to 5.14 mm. This suggests that while low precipitation days saw an increase in rainfall, extreme rainfall events became less frequent compared to 2009. The increased minimum indicates more consistent rainfall, reducing the occurrence of very dry days, but the decrease in maximum rainfall reflects a reduction in very intense rain events. By 2015, the maximum rainfall reached its peak for the entire period at 7.01 mm, while the minimum value slightly decreased to 1.34 mm (Figure 7). This record-high maximum rainfall indicates that 2015 experienced some of the most intense rain events in the South-South zone over the two-decade span. Such extreme rainfall could have led to challenges such as flash floods and infrastructure strain. The lower minimum suggests that while some days had extremely heavy rain, others remained relatively dry, maintaining a wide variability.




Figure 7: Mean Precipitation Across States in the South - south region in 2015

In 2017, the minimum rainfall value was recorded at 1.22 mm, a slight decrease from previous years, while the maximum rainfall reached 6.68 mm. This year is marked by a relatively wide gap between the minimum and maximum values, suggesting both dry and extremely wet days. The high maximum rainfall indicates continued occurrences of significant rain events, while the lower minimum points to periodic dry conditions. In 2020, the minimum rainfall remained stable at 1.23 mm, while the maximum dropped significantly to 5.38 mm. The reduced maximum suggests that extreme rainfall events were less frequent compared to the peak years like 2015. However, the stable minimum value reflects consistent

rainfall on drier days. This could indicate a more moderate rainfall distribution, with fewer extremes, providing a more balanced hydrological cycle in the region for that year.

3.3 Soil Moisture

In 2000, the minimum soil moisture recorded was $0.17 \text{ m}^3/\text{m}^3$, while the maximum value reached $0.46 \text{ m}^3/\text{m}^3$. This range reflects moderate soil moisture conditions, indicating that the region experienced relatively balanced moisture levels (Figure 8). The maximum of 0.46 suggests significant water retention capacity during certain periods, while the minimum of 0.17 suggests that some areas were prone to drying out more quickly.




Figure 8: Mean Soil Moisture Across States in the South - south region in 2000

By 2003, there was an increase in the maximum soil moisture to $0.47 \text{ m}^3/\text{m}^3$, while the minimum remained steady at $0.17 \text{ m}^3/\text{m}^3$. The increase in the maximum value suggests slightly higher moisture retention compared to 2000, indicating that certain regions received more substantial precipitation or retained moisture better during this period. In 2006, the minimum soil moisture rose slightly to $0.174 \text{ m}^3/\text{m}^3$, while the maximum remained constant at $0.47 \text{ m}^3/\text{m}^3$. The increase in the minimum suggests more consistent moisture availability across the region, signaling less variability in soil drying and moisture retention compared to earlier years. The year 2009 saw an increase in the minimum soil moisture to $0.18 \text{ m}^3/\text{m}^3$, while the maximum remained stable at $0.47 \text{ m}^3/\text{m}^3$. The rise in the minimum moisture value indicates that the soil across the South-South geopolitical zone retained higher moisture levels throughout the year. This year shows relatively high overall moisture content, with both minimum and maximum values being among the highest observed. Similar to 2009, 2012 maintained a minimum soil moisture of $0.18 \text{ m}^3/\text{m}^3$ and a maximum of $0.47 \text{ m}^3/\text{m}^3$. The consistency in both minimum and maximum values between 2009 and 2012 suggests stable

moisture conditions during this period, indicating no significant fluctuations in soil moisture retention across the region. In 2015, the minimum soil moisture remained at $0.18 \text{ m}^3/\text{m}^3$, but the maximum dropped slightly to $0.45 \text{ m}^3/\text{m}^3$ (Figure 9). The decrease in maximum soil moisture may indicate a reduction in overall precipitation or changes in soil properties, reducing the ability of certain areas to retain as much water as in previous years.




Figure 9: Mean Soil Moisture Across States in the South - south region in 2015

The year 2017 saw the minimum soil moisture decrease slightly to $0.17 \text{ m}^3/\text{m}^3$, while the maximum remained steady at $0.46 \text{ m}^3/\text{m}^3$. This minor reduction in minimum moisture reflects a slight increase in variability, with some areas experiencing slightly drier conditions compared to the previous years. In 2020, the minimum soil moisture further decreased to $0.16 \text{ m}^3/\text{m}^3$, while the maximum also dropped to $0.45 \text{ m}^3/\text{m}^3$. The decrease in both minimum and maximum values suggests drier conditions across the region compared to earlier years, indicating that soil moisture retention may have been less efficient or that the area received less precipitation during this period.

The comprehensive the soil moisture data for the South-South region of Nigeria from 2000 to 2020 is presented in Table 1. Table 1 demonstrates fluctuations in minimum, maximum, mean, and standard deviation (SD) values, reflecting the interplay of climatic and environmental factors. These variations highlight the sensitivity of soil moisture to changes in precipitation, temperature, and land-use practices, which have significant implications for agriculture, water resource management, and ecosystem stability.

Table 1: Variation in soil moisture (m^3/m^3) in South-South Region of Nigeria from 2000 to 2020

Year	Min	Max	Mean	SD
2000	0.17	0.46	0.315	0.084
2003	0.17	0.47	0.320	0.087
2006	0.174	0.47	0.322	0.085
2009	0.18	0.47	0.325	0.084
2012	0.18	0.47	0.325	0.084
2015	0.18	0.45	0.315	0.078
2017	0.17	0.46	0.315	0.084
2020	0.16	0.45	0.305	0.084

Researcher's analysis 2024

Trend Analysis

The minimum soil moisture values range from $0.16 \text{ m}^3/\text{m}^3$ (2020) to $0.18 \text{ m}^3/\text{m}^3$ (2009, 2012, and 2015). The decline to $0.16 \text{ m}^3/\text{m}^3$ in 2020 indicates drier conditions in some areas, potentially due to reduced precipitation or increased evaporation rates. The relatively higher minimum values in earlier years suggest more consistent baseline moisture levels, which could support vegetation during dry periods. Maximum soil moisture values range from $0.45 \text{ m}^3/\text{m}^3$ (2015 and 2020) to $0.47 \text{ m}^3/\text{m}^3$ (2003, 2006, 2009, and 2012). The slight decline in maximum values after 2012 suggests reduced soil water retention capacity, possibly due to changes in rainfall patterns, soil degradation, or increased land use intensity. The mean soil moisture shows a slight increase from $0.315 \text{ m}^3/\text{m}^3$ (2000) to a peak of $0.325 \text{ m}^3/\text{m}^3$ (2009 and 2012), followed by a gradual decline to $0.305 \text{ m}^3/\text{m}^3$ in 2020. The decline in mean soil moisture after 2012 suggests drier overall conditions, likely driven by a combination of lower precipitation and higher temperatures. The SD values remain relatively stable, ranging from 0.078 (2015) to 0.087 (2003), indicating consistent variability in soil moisture levels over the years. The lower SD in 2015 reflects less variability in soil moisture, potentially due to uniform rainfall distribution or homogeneous soil conditions during that year.

Similarly, the comprehensive precipitation data for the South-South region of Nigeria, over two decades (2000–2020) reveals significant variability in daily rainfall patterns is presented in Table 2. These variations highlight the complexity of rainfall distribution and its potential implications for agriculture, water resource management, and ecosystem stability in the region. The minimum daily precipitation fluctuates between 1.04 mm (2003) and 1.55 mm (2012). This variation indicates that the region experienced dry days across all years, with 2003 being the driest. The relatively higher minimum in 2012 suggests fewer extremely dry days, which could have been influenced by increased baseline moisture levels or consistent rainfall. The maximum precipitation ranges from 5.14 mm (2012) to 7.01 mm (2015), showing a steady increase in peak rainfall events up until 2015. The peak in 2015 suggests that extreme rain events became more pronounced, potentially linked to localized heavy storms or climate-driven anomalies. A decline in maximum precipitation in subsequent years, such as 2020 (5.38 mm), reflects less intense rainfall events during those periods. The mean daily precipitation values range from 3.13 mm (2003) to 4.18 mm (2015), revealing variations in average rainfall distribution. A dip in the mean in 2003 indicates lower overall rainfall, while the peak in 2015 highlights a year of significantly wetter conditions, possibly tied to increased storm activity or prolonged rainy seasons. The SD, which measures the variability of daily precipitation, fluctuates from 0.90 mm (2012) to 1.42 mm (2015). A higher SD, particularly in 2015, reflects a broader range of rainfall intensities, including more

extreme weather events. In contrast, a lower SD in 2012 indicates more consistent daily rainfall, with fewer extreme variations.

Table 2: Variation in, Precipitation (mm/day) in South-South Region of Nigeria from 2000 to 2020

Year	Min	Max	Mean	SD
2000	1.45	5.39	3.42	0.98
2003	1.04	5.22	3.13	1.05
2006	1.32	5.82	3.57	1.12
2009	1.35	6.88	4.12	1.38
2012	1.55	5.14	3.34	0.90
2015	1.34	7.01	4.18	1.42
2017	1.22	6.68	3.95	1.36
2020	1.23	5.38	3.31	1.05

Researcher's analysis 2024

3.4 The Role of Soil Moisture in Moderating Vegetation Response to Climate

Soil moisture serves as a vital regulator in the interaction between climatic variables and vegetation dynamics. In the South-South region of Nigeria, this relationship is particularly critical due to the region's reliance on agriculture and the prevalence of climate variability. Soil moisture acts as a buffer, mitigating the impacts of extreme weather conditions by maintaining water availability for vegetation, even during periods of low precipitation. This capacity underscores its importance in sustaining plant growth and ecosystem health in the face of climatic fluctuations. The findings in this study indicate that soil moisture levels are closely linked to precipitation and temperature trends, with significant implications for vegetation responses. For instance, during periods of high rainfall, soil moisture retention supports robust vegetation growth by providing essential hydration for root systems. Conversely, during drier conditions or extended periods of high temperatures, reduced soil moisture levels can exacerbate vegetation stress, leading to declines in productivity and biodiversity. The temporal trends observed in the region highlight that shifts in soil moisture availability are a critical factor influencing vegetation resilience to climate variability. Seasonal variations in soil moisture further illustrate its moderating role. During the wet season, the increased soil water content facilitates rapid vegetation growth and higher photosynthetic activity. In contrast, the dry season exposes plants to water stress, particularly in areas where soil moisture retention is low.

This dynamic underscores the need for effective land management practices that enhance soil moisture retention, such as mulching, agroforestry, and improved irrigation techniques. These strategies can help maintain vegetation productivity, even under adverse climatic conditions. The coupling between soil moisture and atmospheric conditions also creates feedback loops that influence vegetation responses. For example, elevated soil moisture levels enhance local evaporation, contributing to cloud formation and potentially increasing rainfall. This self-reinforcing mechanism can stabilize vegetation dynamics during transitional climatic phases. However, prolonged deviations in soil moisture—either excess or deficit—can disrupt this balance, leading to adverse ecological outcomes, such as soil erosion, reduced plant cover, and habitat loss. The study also emphasizes the role of soil moisture in moderating extreme climatic events. During floods, adequate soil moisture capacity can reduce runoff and prevent erosion, protecting vegetation cover. Similarly, during droughts, residual soil moisture can sustain vegetation for longer periods, mitigating the immediate impacts of water scarcity. These functions are critical for maintaining ecosystem stability in the South-South region,

where climatic extremes are becoming more frequent due to global climate change. In conclusion, soil moisture is a cornerstone in the complex interplay between climate and vegetation in the South-South region of Nigeria. Its ability to regulate water availability, support vegetation growth, and influence local climatic feedback mechanisms makes it indispensable for sustaining agricultural productivity and ecological resilience. Future conservation and land management initiatives must prioritize strategies that enhance soil moisture retention and mitigate the adverse impacts of climate variability on vegetation.

4.0 CONCLUSION

The analysis of climate variability and soil moisture dynamics in the South-South region of Nigeria underscores the profound impact of changing climatic patterns on ecosystems, agriculture, and water resources. Increasing temperatures, fluctuating precipitation, and declining soil moisture trends collectively pose significant challenges to the region's environmental and socioeconomic sustainability. To address these issues effectively, adaptive strategies and resource management practices must be prioritized to enhance resilience and ensure sustainable development.

Recommendations

- i. Implement sustainable agricultural practices, such as conservation tillage, mulching, and the use of cover crops, to improve soil structure and moisture retention.
- ii. Promote agroforestry systems that integrate trees and crops to maintain soil stability and reduce water loss through evapotranspiration.
- iii. Develop irrigation systems tailored to local climatic conditions, including the adoption of drip irrigation and rainwater harvesting to optimize water use efficiency.
- iv. Encourage the cultivation of drought-tolerant and water-efficient crop varieties to mitigate the impacts of irregular rainfall and prolonged dry spells.
- v. Provide training and support for farmers to adopt climate-smart agricultural practices, including soil and water conservation techniques.

By implementing these recommendations, the South-South region of Nigeria can better adapt to the challenges posed by climate variability while safeguarding its natural resources and supporting its agricultural and ecological systems. A coordinated effort among stakeholders will be essential to achieve sustainable development and resilience in the face of a changing climate.

REFERENCES

- Adeogun, A. G., & Adeogun, T. R. (2014). GIS-based analysis of hydrological parameters in Nigeria. *Journal of Environmental Management*, 30(3), 211–225.
- Adeogun, A. G., et al. (2023). Application of hydrological modeling and GIS in soil moisture studies. *Journal of Hydrology Research*, 57(4), 298–310.
- Adesina, A. A., et al. (2020). The environmental impacts of oil exploration in Nigeria. *Energy and Environment Studies*, 45(3), 12–24.
- Agbonaye, P., & Izinyon, O. (2021). Performance of satellite-based rainfall products in Nigeria. *African Journal of Climate Research*, 12(1), 112–124.
- Ardilouze, C., et al. (2018). Feedback between soil moisture and precipitation in tropical ecosystems. *Climate Dynamics*, 52(1), 235–248.
- Berg, A., et al. (2014). Climate feedbacks and soil moisture interactions. *Nature Climate Change*, 4(7), 501–505.
- Carón, L., et al. (2014). Tropical precipitation dynamics influenced by soil moisture anomalies. *Journal of Climate*, 27(4), 12–23.
- Das, R., & Maity, P. (2014). Linking soil moisture variability to crop yield in tropical regions. *Agricultural Water Management*, 120(3), 42–54.
- Dembélé, M., et al. (2022). Climate variability and precipitation modeling using GCMs. *Climatic Change Research*, 89(2), 134–147.
- Dirmeyer, P. A., & Norton, S. S. (2018). Soil moisture feedback on climate systems. *Journal of Climate Dynamics*, 54(3), 41–53.
- Ekness, P., & Randhir, T. O. (2015). Hydrological responses to soil moisture variability. *Water Resource Research*, 56(4), 567–582.
- Enuoh, O. O., & Ogogo, U. (2018). Mangrove conservation and local livelihoods in the Niger Delta. *Journal of Environmental Studies*, 45(3), 210–224.
- Fernandez-Long, M., et al. (2020). Soil moisture and hydrological cycle dynamics. *Journal of Hydrology and Climate Change*, 13(2), 201–223.
- Ghajarnia, N., et al. (2020). Monitoring soil moisture changes in tropical regions. *Environmental Monitoring*, 58(2), 120–139.
- Guo, X., et al. (2016). Using satellite data to track soil moisture variability. *Journal of Remote Sensing Applications*, 34(1), 109–125.
- Han, Z., et al. (2020). The relationship between soil moisture and crop yield in tropical climates. *Agricultural and Forest Meteorology*, 79(3), 123–136.
- Huang, Y., et al. (2022). Advancing soil moisture monitoring through remote sensing. *Advances in Climate Data*, 46(2), 78–98.
- Krakauer, N. Y., et al. (2010). Soil moisture dynamics in tropical and arid regions. *Water Resources Research*, 22(4), 333–345.
- Li, H., & Rodell, M. (2013). Modeling soil moisture variations under changing climatic conditions. *Climate Change Journal*, 65(3), 45–57.

- Li, Z., et al. (2019). Soil moisture anomalies and agricultural impacts. *Journal of Hydrology*, 78(2), 234–246.
- Liu, Y. (2023). Trends in soil moisture variability under climate change. *Journal of Environmental Studies*, 56(1), 12–27.
- Lorenz, C., et al. (2016). Seasonal dynamics of soil moisture and vegetation in tropical regions. *Journal of Climate Research*, 28(3), 77–89.
- Merem, E. C., et al. (2019). Rainfall variability and agricultural impacts in Nigeria. *Journal of Agricultural Sciences*, 18(2), 145–162.
- Mittelbach, H., & Seneviratne, S. I. (2012). Soil moisture heterogeneity and vegetation cover effects. *Journal of Climate Dynamics*, 38(1), 23–37.
- Morrison, M. R., & Wagner, B. (2020). Soil moisture changes and groundwater recharge in tropical climates. *Hydrological Science Bulletin*, 67(3), 167–190.
- Myeni, L., et al. (2021). Effects of vegetation cover on soil moisture retention. *Journal of Environmental Monitoring*, 45(2), 301–318.
- Ogbu, C., et al. (2020). Evaluation of CHIRPS data in Nigeria. *Climate Data Journal*, 5(1), 67–78.
- Odusanya, A., et al. (2019). Satellite-based evapotranspiration data for hydrological modeling. *Journal of Hydrology*, 78(3), 109–127.
- Okeola, O., et al. (2023). Advances in soil moisture measurement techniques. *Journal of Environmental Hydrology*, 29(4), 56–68.
- Roux, P., et al. (2013). Soil moisture influences on tropical rainfall patterns. *Environmental Science Journal*, 47(2), 81–96.
- Saeed, F., et al. (2013). Soil moisture and crop yield variability. *Agricultural Water Management*, 120(1), 23–33.
- Salley, M. A., et al. (2016). Feedback effects between soil moisture and precipitation. *Journal of Hydrometeorology*, 17(2), 278–289.
- Schwingshackl, C., et al. (2018). Variability of soil moisture across tropical ecosystems. *Journal of Hydrological Research*, 54(2), 134–148.
- Senanayake, I., et al. (2022). Impacts of soil moisture anomalies on seasonal weather. *Journal of Tropical Meteorology*, 48(3), 123–139.
- Si, L., et al. (2023). Soil moisture variability and implications under future climate scenarios. *Journal of Climate Variability*, 29(3), 201–225.
- Song, J., et al. (2019). Soil moisture monitoring using satellite-based data. *Advances in Climate Data Science*, 22(4), 456–470.
- Song, J., et al. (2021). Advances in remote sensing technologies for soil moisture. *Journal of Geophysical Monitoring*, 34(2), 78–98.
- Tinubu, S., et al. (2019). Challenges of climate variability in Nigeria's agriculture. *African Environmental Studies*, 7(2), 99–108.

- Uguru H., et al. (2022). Comprehensive assessment of the effect of various anthropogenic activities on the groundwater quality. *Sci. Adv. Mater.*, 14: 462–474
- Uguru, H., et al. (2024). Exposure to airborne pollutants in urban and rural areas: levels of metals and microorganisms in PM10 and gaseous pollutants in ambient air. *Air Quality, Atmosphere & Health*.
- Urban, G., et al. (2015). Soil moisture and hydrological cycle feedbacks in tropical regions. *Journal of Climate Dynamics*, 21(1), 145–166.
- Usman, A., et al. (2018). Analysis of extreme precipitation trends in Nigeria. *Journal of Hydrology and Climate Change*, 13(3), 185–198.
- Vezhapparambu, S., et al. (2019). Rainfall and soil moisture variability in tropical regions. *Journal of Climate Research*, 32(1), 109–125.
- Wang, X., et al. (2018). Regional impacts of soil moisture on tropical hydrological systems. *Journal of Climate Research*, 54(2), 201–222.
- Wing, B., et al. (2016). Coupling between soil moisture and atmospheric systems. *Hydrology Journal*, 37(3), 89–101.
- Zhang, T., et al. (2011). Feedbacks between soil moisture and climate systems. *Journal of Hydrology*, 52(4), 321–338.
- Zhu, Y., et al. (2019). Soil moisture changes under global warming. *Journal of Environmental Hydrology*, 45(1), 65–89