IJAAR PUBLISHING®
Scholarly Journal of Science and Technology Research & Development | ISSN: 2955-0807
Vol. 4, Issue 3 (2025) | www.ijaar.org/sjstrd

BIODEGRADATION: A FORMIDABLE APPROACH TO CURBING OIL SPILLAGE

Okatahi Aminu Mohammed[1]; Enejoh Omejeh[1]; Ahmad Abdulrazaq Itopa[1]

[1]Kogi State Polytechnic, Lokoja, Kogi State

Abstract

Oil spills pose severe environmental threats, contaminating marine and terrestrial ecosystems and causing long-term ecological and economic damage. Biodegradation, the natural process by which microorganisms break down oil hydrocarbons into simpler compounds, offers a sustainable and eco-friendly approach to mitigate these disasters. This review explores the mechanisms of biodegradation, its advantages over traditional remediation methods, current challenges, and emerging biotechnological advancements that enhance its efficacy in curbing oil spillage.

Keywords: biodegradation; terrestrial; ecosystem; hydrocarbons; eco-friendly

Mohammed, O. A., Omejeh, E., & Itopa, A. A. (2025). Biodegradation: A Formidable Approach to Curbing Oil Spillage. *Scholarly Journal of Science and Technology Research & Development. 4*(3), 25-36. DOI: https://doi.org/10.5281/zenodo.16459309

1.0 Introduction

Oil spills are among the most devastating environmental disasters, causing significant ecological and economic damage to marine and terrestrial ecosystems. The release of crude oil into the environment, whether through accidents, operational failures, or illegal discharges, poses severe threats to biodiversity, water quality, and human health.

1.1 Definition of Oil Spillage

Oil spillage refers to the accidental or intentional release of petroleum hydrocarbons into the environment, particularly marine and terrestrial ecosystems, causing significant ecological and economic damage (Atlas & Hazen, 2011).

Traditional methods of oil spill remediation, such as physical removal (e.g., skimming, booms) and chemical dispersants, are often costly, labor-intensive, and may introduce secondary pollutants (Atlas & Hazen, 2011). In response to these challenges, biodegradation has emerged as a promising, eco-friendly approach to mitigate oil spillage. Biodegradation leverages the natural metabolic capabilities of microorganisms to break down and mineralize hydrocarbons in oil, transforming them into less harmful compounds like carbon dioxide, water, and biomass. This paper provides a comprehensive introduction to biodegradation, highlighting its mechanisms, advantages, limitations, and its role as a sustainable strategy for addressing oil spills.

Biodegradation is the natural process by which microorganisms, such as bacteria, fungi, and algae, degrade organic pollutants, including petroleum hydrocarbons, into simpler, non-toxic substances. In the context of oil spills, biodegradation involves the microbial breakdown of complex hydrocarbons—such as alkanes, aromatics, and polycyclic aromatic hydrocarbons (PAHs)—into carbon dioxide, water, and microbial biomass (Das & Chandran, 2011). This process is facilitated by enzymes produced by microorganisms that catalyze the oxidation of hydrocarbons, enabling microbes to use oil as a carbon and energy source.

The efficiency of biodegradation depends on several factors, including the type of oil, environmental conditions (e.g., temperature, pH, oxygen availability), and the presence of suitable microbial communities. Crude oil is a complex mixture of hydrocarbons, including alkanes (straight and branched), cycloalkanes, and aromatic compounds, each with varying susceptibility to microbial degradation. For instance, lighter hydrocarbons (e.g., n-alkanes) are more readily biodegradable, while heavier compounds like PAHs are more resistant due to their complex molecular structures (Leahy & Colwell, 1990).

1.2 Mechanisms of Biodegradation

Biodegradation of oil occurs through aerobic and anaerobic pathways, with aerobic degradation being the most efficient and commonly observed in oil spill scenarios. Under aerobic conditions, microorganisms use oxygen as an electron acceptor to oxidize hydrocarbons, producing energy and breaking down the oil into simpler compounds. The process begins with the activation of hydrocarbons by oxygenase enzymes, which introduce oxygen atoms into the hydrocarbon structure, making it more amenable to further degradation (Rojo, 2009). Key microbial genera involved in aerobic oil degradation include Pseudomonas, Alcanivorax, and Rhodococcus, which are known for their hydrocarbon-degrading capabilities (Figure 1).

Anaerobic biodegradation, while slower, occurs in oxygen-limited environments, such as deep-sea sediments or waterlogged soils. Anaerobic microbes use alternative electron acceptors, such as nitrate, sulfate, or iron, to metabolize hydrocarbons. Although less

efficient, anaerobic degradation is critical in environments where oxygen is scarce, such as in deep marine ecosystems affected by oil spills like the Deepwater Horizon incident (Hazen et al., 2010).




Figure 1: Diagram of aerobic biodegradation of hydrocarbons. (Fritsche and Hofrichter 2000)

Generally, the biodegradation, that occur very fast and completely for the majority of organic contaminants, was achieved in the aerobic restraint. Figure 1 exhibits the major precept of the aerobic biodegradation of hydrocarbon substances (Fritsche and Hofrichter 2000). The first intracellular crack of organic contaminants in the environment was via an oxidation operation. The environmental paths of biodegradation of hydrocarbons transform such organic contaminants gradually to fragments through the centrally mediated metabolism such as the cycle of tricarboxylic acid. The biological synthesis of the cell mass occurs, primarily, from the metabolism of the centric precursors, for example, acetyl-CoA, pyruvate, and succinate. In addition, sugars, needed in different biological synthesis, are mainly synthesized by gluconeogenesis process (Fritsche and Hofrichter 2000).

The effectiveness of biodegradation in remediating oil spills is governed by several environmental and biological factors, as summarized in Table 1. These factors include: Oil

Composition, Nutrient Availability, Temperature, Oxygen Availability, And Microbial Community:

Table 1: Key Factors Influencing Biodegradation of Oil Spills

Factor	Impact on Biodegradation	Example Environment
Oil Composition	Lighter hydrocarbons degrade faster; PAHs are slower	Crude oil vs. heavy fuel oil
Nutrient Availability	Nitrogen and phosphorus enhance microbial growth	Nutrient-poor marine waters
Temperature	Higher temperatures increase degradation rates	Tropical vs. Arctic environments
Oxygen Availability	Aerobic degradation is faster than anaerobic	Surface waters vs. deep-sea sediments
Microbial Community	Hydrocarbon-degrading microbes enhance efficiency	Coastal vs. open ocean ecosystems

(Source: Adapted from Leahy & Colwell, 1990; Atlas & Hazen, 2011)

2.0 What is Biodegradation?

Biodegradation is the microbial-mediated process of breaking down complex organic compounds, such as the hydrocarbons in crude oil, into simpler substances like carbon dioxide, water, and biomass. Microorganisms, including bacteria, fungi, and algae, utilize hydrocarbons as a carbon and energy source, transforming them through enzymatic processes. This natural remediation method is particularly effective for oil spills because it leverages the metabolic capabilities of indigenous or introduced microorganisms to degrade pollutants.

2.1 Mechanisms of Biodegradation

The biodegradation of oil involves several biochemical pathways:

2.1.1 Aerobic Biodegradation

In the presence of oxygen, microorganisms oxidize hydrocarbons using enzymes like monooxygenases and dioxygenases. For example, alkanes are broken down into alcohols, aldehydes, and fatty acids, which are further metabolized into CO_2 and H_2O. Microorganisms use oxygen as an electron acceptor to oxidize hydrocarbons into carbon dioxide, water, and biomass. For example, bacteria like Pseudomonas and Alcanivorax produce enzymes such as alkane hydroxylases that initiate hydrocarbon breakdown (Rojo, 2009).

2.1.2 Anaerobic Biodegradation

In oxygen-limited environments (e.g., deep-sea sediments), microbes use alternative electron acceptors like nitrate or sulfate. This process is slower but relevant in certain spill scenarios, such as deepwater oil spills (Kimes et al., 2014).

2.1.3 Enzymatic Pathways

Key enzymes, including monooxygenases and dioxygenases, facilitate the breakdown of complex hydrocarbons into simpler intermediates, which are further metabolized via microbial pathways (Das & Chandran, 2011).

Table 2: Key Microorganisms Involved in Oil Biodegradation

Microorganism Type	Examples	Hydrocarbon Preference	Environment
Bacteria	Pseudomonas spp., Alcanivorax spp.	Alkanes, Aromatics	Marine, Soil
Fungi	Aspergillus spp., Penicillium spp.	Polycyclic Aromatic Hydrocarbons (PAHs)	Soil
Algae	Chlorella spp.	Light hydrocarbons	Aquatic

2.2 Factors Influencing Biodegradation

The efficiency of biodegradation depends on several environmental and biological factors:

2.2.1 Hydrocarbon Composition

Lighter hydrocarbons (e.g., alkanes) degrade faster than heavier ones (e.g., PAHs).

2.2.2 Oxygen Availability

Aerobic degradation is faster and more efficient than anaerobic processes.

2.2.3 Nutrient Availability

Nitrogen, phosphorus, and other nutrients are essential for microbial growth and activity.

2.2.4 Temperature

Warmer temperatures (20–40°C) enhance microbial activity, while extreme temperatures slow degradation.

2.2.5 pH and Salinity

Most oil-degrading microbes thrive in neutral pH and moderate salinity.

2.2.6 Microbial Population

The presence of hydrocarbon-degrading microbes, either indigenous or introduced, is critical (Hazen, T. C. et al., 2010).

2.3 Applications of Biodegradation in Oil Spill Remediation

Biodegradation is applied through two main strategies:

2.3.1 Natural Attenuation

Relies on indigenous microorganisms to naturally degrade oil without human intervention. This is effective in environments with favorable conditions but can be slow.

2.3.2 Bioremediation

Introduces specialized hydrocarbon-degrading microbes to enhance degradation rates (Bioaugmentation). Adds nutrients (e.g., nitrogen, phosphorus) or oxygen to stimulate native microbial activity (Biostimulation). Uses modified microbes with enhanced degradation capabilities, though this is less common due to regulatory concerns (Genetically Engineered Microbes) (Atlas, et al., 2011).

2.4 Advantages of Biodegradation

It produces non-toxic byproducts, minimizing environmental harm. It requires minimal infrastructure compared to physical or chemical cleanup methods. Leverages natural processes, reducing the need for invasive interventions and. Applicable to various environments, including marine, terrestrial, and freshwater systems (Atlas & Hazen, 2011).

3.0 Mechanisms of Biodegradation in Oil Spill Remediation

Oil spills pose significant environmental threats, contaminating marine and terrestrial ecosystems with complex mixtures of hydrocarbons. Biodegradation, the microbial breakdown of organic compounds, offers a sustainable approach to mitigate oil spill impacts by transforming hydrocarbons into less harmful substances like carbon dioxide, water, and biomass. This process relies on the metabolic capabilities of microorganisms such as bacteria, fungi, and algae. Understanding the mechanisms of biodegradation is critical to optimizing its application in oil spill remediation. This paper explores the biochemical pathways, microbial players, environmental factors, and enhancement strategies involved in biodegradation, with emphasis on its role as a formidable tool for oil spill cleanup.

3.1 Biochemical Pathways of Biodegradation

Biodegradation of oil involves the enzymatic breakdown of hydrocarbons, primarily alkanes, alkenes, and polycyclic aromatic hydrocarbons (PAHs). The process occurs through aerobic or anaerobic pathways, depending on oxygen availability.

3.1.1 Aerobic Biodegradation

Aerobic biodegradation is the dominant mechanism in oxygen-rich environments like surface waters or shorelines. Microorganisms use oxygen as an electron acceptor to oxidize hydrocarbons. The key steps include: Enzymes such as alkane hydroxylases or monooxygenases introduce oxygen into hydrocarbon molecules, forming alcohols. For example, alkanes are oxidized to primary alcohols via the enzyme alkane 1-monooxygenase (Hassanshahian & Boroujeni, 2020). Alcohols are further oxidized to aldehydes, then fatty acids, which enter the beta-oxidation pathway, producing acetyl-CoA for the microbial tricarboxylic acid (TCA) cycle (Das & Chandran, 2011). PAHs are degraded via dioxygenases, which incorporate two oxygen atoms into aromatic rings, forming catechols that are cleaved and funneled into central metabolic pathways (Harayama et al., 2004).




Figure 2: Simplified Aerobic Biodegradation Pathway of Alkanes

The degradation pathways of n-alkanes. a, The aerobic degradation pathways of n-alkanes; b the hexadecane degradation pathway of strain G. thermodenitrificans NG80-2 (Feng et al. 2007); c the anaerobic degradation pathways of n-alkanes

3.1.2 Anaerobic Biodegradation

In oxygen-limited environments, such as deep-sea sediments or saturated soils, anaerobic microorganisms use alternative electron acceptors (e.g., nitrate, sulfate, or CO_2). Anaerobic degradation is slower but critical for spills in anoxic zones. Key mechanisms include: Alkanes are activated by adding fumarate, forming alkylsuccinates, which are further metabolized (Widdel & Rabus, 2001). Anaerobic bacteria like sulfate-reducing Desulfobacter species degrade PAHs via carboxylation or methylation, though this process is less efficient than aerobic pathways (Prince et al., 2010).

3.2 Key Microbial Players

Microorganisms capable of degrading hydrocarbons are diverse, with bacteria being the most studied due to their metabolic versatility. Key genera include:

3.2.1 Alcanivorax borkumensis

A marine bacterium specializing in alkane degradation, dominant in oil-contaminated waters (Yakimov et al., 2007).

3.2.2 Pseudomonas spp

Versatile degraders of alkanes and PAHs, producing biosurfactants to enhance oil bioavailability (Das & Chandran, 2011).

3.2.3 Rhodococcus spp

Effective in degrading complex hydrocarbons in soil and cold environments (Atlas, 1995).

3.2.4 Fungi and Algae

Fungi like Aspergillus and algae like Chlorella contribute to PAH degradation, though their role is less prominent (Cerniglia & Sutherland, 2006).

Table 3: Key Microorganisms and Their Roles in Oil Biodegradation

Microorganism	Hydrocarbon Target	Environment	Key Enzymes/Biosurfactants
Alcanivorax borkumensis	Alkanes	Marine	Alkane hydroxylase
Pseudomonas spp.	Alkanes, PAHs	Soil, Water	Monooxygenases, Biosurfactants
Rhodococcus spp.	Alkanes, Aromatics	Soil, Cold Regions	Cytochrome P450
Aspergillus spp.	PAHs	Soil	Ligninolytic enzymes

3.3 Enhancement Strategies for Biodegradation

To overcome limitations, bioremediation strategies enhance microbial activity:

3.3.1 Biostimulation

Adding nutrients (e.g., nitrogen, phosphorus) boosts indigenous microbial populations. During the Deepwater Horizon spill, fertilizers increased degradation rates by 20–40% (Hazen et al., 2010).

3.3.2 Bioaugmentation

Introducing specialized microbial consortia enhances degradation of recalcitrant compounds. Genetically engineered strains, like Pseudomonas with enhanced alkB genes, show promise (Joutey et al., 2013).

3.3.3 Biosurfactants

Microbial or synthetic surfactants increase oil bioavailability by emulsifying hydrocarbons, improving microbial access (Das & Chandran, 2011).

3.3.4 Nanotechnology

Nanoparticle-based nutrient delivery systems improve nutrient dispersal in marine environments (Alabresm et al., 2018).

4.0 Case Studies and Practical Applications

4.1Deepwater Horizon (2010)

Biostimulation with nitrogen and phosphorus fertilizers enhanced Alcanivorax activity, degrading significant portions of oil in the Gulf of Mexico (Hazen et al., 2010). Studies estimate 20–40% of the oil was biodegraded within months.

4.2 Exxon Valdez (1989)

Fertilizer application on Alaskan shorelines increased alkane degradation by native microbes, reducing oil residues (Atlas, 1995).

4.3 Emerging Research

Genetic engineering of Alcanivorax to target PAHs and nanotechnology-based nutrient delivery are advancing bioremediation efficacy (Joutey et al., 2013; Alabresm et al., 2018).

5.0 Challenges and Future Directions

5.1 Challenges

Despite its potential, biodegradation faces challenges:

Biodegradation is slower than mechanical methods, often requiring weeks to months for significant cleanup; Incomplete Degradation: Heavy hydrocarbons like asphaltenes resist degradation, leaving residual pollutants (Prince et al., 2010); Nutrient additions can cause algal blooms, disrupting ecosystems (Hazen et al., 2010) and; Low temperatures in Arctic spills limit microbial activity, necessitating cold-adapted microbes (Hassanshahian & Boroujeni, 2020).

Table 4: Comparison of Biodegradation with Other Oil Spill Remediation Methods

Method	Pros	Cons	Cost
Biodegradation	Eco-friendly, sustainable	Slow, site-specific	Low
Chemical Dispersants	Fast, effective in water	Toxic residues, environmental harm	High
Mechanical Removal	Immediate, visible results	Labor-intensive, incomplete	High

5.2 Future Directions

The future direction include: Engineering microbes with enhanced degradation genes (e.g., alkB, nahAc) for faster and broader hydrocarbon breakdown; Combining biodegradation with dispersants or physical methods for comprehensive cleanup and; Developing psychrophilic (cold-tolerant) microbes for Arctic spills.

6.0 Conclusion

Biodegradation is a formidable approach to oil spill remediation, leveraging microbial metabolic pathways to degrade hydrocarbons in offering a sustainable, cost-effective, and environmentally friendly alternative to traditional cleanup methods. Aerobic and anaerobic mechanisms, driven by key players like Alcanivorax and Pseudomonas, are modulated by environmental factors and enhanced through biostimulation, bioaugmentation, and emerging technologies. While challenges like slow degradation rates and incomplete breakdown

persist, advancements in synthetic biology and nanotechnology promise to enhance its efficacy. Ongoing research into microbial genomics and bioremediation technologies will further enhance the efficacy of biodegradation, making it a cornerstone of oil spill response strategies in the future. By understanding and optimizing these mechanisms, biodegradation can play a central role in sustainable oil spill management.

References

Alabresm, A., Chen, Y. P., Decho, A. W., & Lead, J. (2018). Engineered nanoparticles enhance bioremediation of oil spills in marine environments. Environmental Science & Technology, 52(18), 10326–10334. https://doi.org/10.1021/acs.est.8b02766

Atlas, R. M. (1995). Petroleum biodegradation and oil spill bioremediation. Marine Pollution Bulletin, 31(4–12), 178–182. https://doi.org/10.1016/0025-326X(95)00113-2

Atlas, R. M., & Hazen, T. C. (2011). Oil biodegradation and bioremediation: A tale of the two worst spills in U.S. history. Environmental Science & Technology, 45(16), 6709–6715. https://doi.org/10.1021/es2013227

Cerniglia, C. E., & Sutherland, J. B. (2006). Bioremediation of polycyclic aromatic hydrocarbons by ligninolytic and non-ligninolytic fungi. In B. K. Singh (Ed.), Fungi in bioremediation (pp. 136–187). Cambridge University Press.

Das, N., & Chandran, P. (2011). Microbial degradation of petroleum hydrocarbon contaminants: An overview. Biotechnology Research International, 2011, Article ID 941810. https://doi.org/10.4061/2011/941810

Feng, L., Wang, W., Cheng, J., Ren, Y., Zhao, G., Gao, C., ... & Wang, L. (2007). Genome and proteome of long-chain alkane degrading Geobacillus thermodenitrificans NG80-2 isolated from a deep-subsurface oil reservoir. *Proceedings of the national academy of sciences*, *104*(13), 5602-5607.

Fritsche W, Hofrichter M (2000) Aerobic degradation by microorganisms. In: Klein J (ed) Environmental processes-soil decontamination. Wiley-VCH, Weinheim, pp146–155

Harayama, S., Kasai, Y., & Hara, A. (2004). Microbial communities in oil-contaminated seawater. Current Opinion in Biotechnology, 15(3), 205–214. https://doi.org/10.1016/j.copbio.2004.04.002

Hassanshahian, M., & Boroujeni, N. A. (2020). Biodegradation of hydrocarbons by extremophilic microorganisms. In A. K. Mishra (Ed.), Extremozymes and their industrial applications (pp. 167–189). Academic Press.

Hazen, T. C., Dubinsky, E. A., DeSantis, T. Z., Andersen, G. L., Piceno, Y. M., Singh, N., & Mason, O. U. (2010). Deep-sea oil plume enriches indigenous oil-degrading bacteria. Science, 330(6001), 204–208. https://doi.org/10.1126/science.1195979

Joutey, N. T., Bahafid, W., Sayel, H., & El Ghachtouli, N. (2013). Biodegradation: Involved microorganisms and genetically engineered microorganisms. In Biodegradation - Life of science (pp. 289–320). InTech.

Kimes, N. E., Callaghan, A. V., Suflita, J. M., & Morris, P. J. (2014). Microbial transformation of the Deepwater Horizon oil spill—past, present, and future perspectives. Frontiers in Microbiology, 5, 603. https://doi.org/10.3389/fmicb.2014.00603

Leahy, J. G., & Colwell, R. R. (1990). Microbial degradation of hydrocarbons in the environment. Microbiological Reviews, 54(3), 305–315. https://doi.org/10.1128/mr.54.3.305-315.1990

Prince, R. C. (2018). Petroleum spill bioremediation in marine environments. Critical Reviews in Environmental Science and Technology, 48(5), 345–367. https://doi.org/10.1080/10643389.2018.1439654

Prince, R. C., Gramain, A., & McGenity, T. J. (2010). Prokaryotic hydrocarbon degraders. In K. N. Timmis (Ed.), Handbook of hydrocarbon and lipid microbiology (pp. 1665–1692). Springer.

Rojo, F. (2009). Degradation of alkanes by bacteria. Environmental Microbiology, 11(10), 2477-2490. https://doi.org/10.1111/j.1462-2920.2009.01948.x

Widdel, F., & Rabus, R. (2001). Anaerobic biodegradation of saturated and aromatic hydrocarbons. Current Opinion in Biotechnology, 12(3), 259–276. https://doi.org/10.1016/S0958-1669(00)00209-3

Yakimov, M. M., Timmis, K. N., & Golyshin, P. N. (2007). Obligate oil-degrading marine bacteria. Current Opinion in Biotechnology, 18(3), 257–266. https://doi.org/10.1016/j.copbio.2007.04.006